

Video Education to Promote Skin Cancer Awareness and **Identification in Spanish**speaking Patients

ABSTRACT

Objective: We sought to evaluate the efficacy of a Spanish-language educational video in teaching primary Spanish speaking patients to recognize benign and malignant lesions and to increase their awareness about skin cancer. Materials and Methods: Thirty-seven subjects were enrolled in study. An instructional video was developed to increase knowledge of benign and malignant lesions, skin cancer awareness, and prevention among Spanish-speaking patients. Two examples each of six common skin lesions (e.g., malignant melanoma, cherry angioma, seborrheic keratosis, benign melanocytic nevus, basal cell carcinoma, and squamous cell carcinoma) were presented as high-quality images to the participants before and after watching the two-minute educational video. A pre- and postvideo survey was used to assess competency. **Results:** The prevideo baseline median score was six points (interquartile range [IQR]: 5–6 points); postviewing median score improved to 11 points (IQR: 11-12 points), which was statistically significant (Median=5 points, IQR: 4–6 points; p<.001). The ability of the participants to identify nonmelanoma skin cancers improved from 74 percent to 98 percent and from 35 percent to 99 percent for squamous cell carcinoma and basal cell carcinoma. Initially, only 30 percent of participants could identify melanoma prior to viewing the video. Afterwards, 97 percent of participants could identify this malignancy. However, the video format preferences were not statistically significant: 67.6 percent of the participants preferred the video format. Conclusion: These results suggest that this educational video is an effective and valuable method to enhance knowledge about skin health and improve identification of skin cancer among Spanish-speaking patients.

KEYWORDS: Dermatology Spanish-speaking, video education, skin cancer education

by DANTE GARCIA, MD; ITISHA S. JEFFERSON, BS; PABLO RAMIREZ, BS; ANGELINA PALOMINO, BS; WILLIAM ADAMS, PhD; JESSICA VERA, BSN; RUBY DE LA TORRE, APN; KRISTIN LEE, MD; ASHLEY ELSENSOHN, MD, MPH; HANA KAZBOUR, MD; and REBECCA TUNG, MD

Dr. Garcia is with the Department of Internal Medicine at Loma Linda University in Loma Linda, California. Mr. Ramirez is with San Juan Bautista School of Medicine in Caguas, Puerto Rico. Dr. Adams is with Biostatistics Core at Stritch School of Medicine of Loyola University Chicago. Ms. Vera is with the Department of Dermatology at Northwestern University in Chicago, Illinois. Dr. Lee and Mses. Jefferson, Palomino, and De La Torre are with the Division of Dermatology at Stritch School of Medicine of Loyola University Chicago in Chicago, Illinois. Dr. Elsensohn is with the Department of Dermatology at the University of California Irvine in Irvine, California. Dr. Kazbour is with the Department of Internal Medicine at Loyola University Chicago Stritch School of Medicine. Dr. Tung is with Florida Dermatology and Skin Cancer Centers in Winter Haven, Florida.

J Clin Aesthet Dermatol. 2020;13(1):41-43

Hispanics are currently the fastest growing ethnic group in the United States (US), comprising nearly a quarter (24.4%) of the population by 2050.1 Although malignant skin lesions are less prevalent among Hispanic individuals, when skin cancer occurs, there is often greater morbidity and mortality in this group when compared to non-Hispanic Caucasian people.^{1–3} Even though Hispanic people frequently present with more advanced stages of skin cancer, they are less likely to implement methods of skin cancer prevention, such as sunscreen use, due to the perception that having a darker skin type confers a lower risk for skin cancer development.4 Video education has been shown to be an effective method for informing individuals on skin cancer prevention and identification.^{5,7} However, there is a lack of multimedia education on this topic that specifically targets Spanish-speaking patients.^{3,5-7} Given

the dearth of accessible Spanish-language educational materials and the need to inform Hispanic communities about skin cancer, we conducted a study to assess the effectiveness of an accessible Spanish-language educational video to increase awareness of skin cancer and improve recognition of benign and malignant lesions in primary Spanish-speaking patients.

METHODS

This study was approved by the Loyola University Chicago Institutional Review Board. Participation by subjects was strictly voluntary. All interested primary Spanish-speaking patients at a family medicine clinic (N=37)in Maywood, Illinois, were enrolled in this study. An anonymous intake form elicited basic patient data, including patient demographics, skin cancer history, full-body skin examination history, typical sunscreen usage, and knowledge about the visual characteristics

FUNDING: No funding was provided.

DISCLOSURES: The authors report no conflicts of interest related to the content of this article.

CORRESPONDENCE: Itisha S. Jefferson; Email: ijefferson@luc.edu

		MEDIAN (IOD)			
DEMOGRAPHICS	N (%)	MEDIAN (IQR)			<i>P</i> -VALUE
		PRE-VIDEO	POST-VIDEO	∆ SCORE	
Sex					0.63
Male	11 (30)	6 (5–6)	11 (10–11)	5 (4-6)	
Female	26 (70)	6 (5–7)	11 (11–12)	6 (4–6)	
Education					
None through Middle School	32 (86)	6 (5–7)	11 (11–12)	5 (4-6)	
High School or College	5 (14)	5 (5–5)	11 (11–12)	6 (6–7)	
Preferred Educational Format					
Written	12 (32)	5 (5-6)	11 (11–12)	6 (5-7)	
Video	25 (68)	6 (5–6)	11 (11–12)	5 (4–6)	
Fitzpatrick Skin Type					
Types I — III	11 (30)	5 (4–6)	11 (11–12)	6 (5-7)	
Types IV — VI	26 (70)	6 (5–7)	11 (11–11)	5 (4-6)	
Outdoor Sunscreen Use					0.005
Never	26 (70)	6 (5–7)	11 (11–11)	5 (4-6)	
Rarely	5 (14)	6 (5–8)	11 (11–11)	5 (2-6)	
Sometimes	6 (16)	5 (5-5)	12 (11–12)	7 (6–7)	
Daily Sunscreen					0.59
No	34 (92)	6 (5–6)	11 (11–12)	5 (4-6)	
Yes	3 (8.1)	5 (5–7)	11 (11–12)	6 (4–7)	
Reapplication of Sunscreen					
No	35 (95)	6 (5–7)	11 (11–12)	5 (4-6)	
Yes	2 (5.4)	5 (5-5)	11 (11–11)	6 (6–6)	

Mean age was 50.92 years (Standard deviation [SD]=10.68)

IQR=Interquartile range.

Statistical significance (p) for the change score among groups was computed using an exact version of the Wilcoxon rank sum test; an exact version of the Kruskal-Wallis test was used for outdoor sunscreen use.

TABLE 2. Pre and post-test percent correct for skin cancer identification.							
SKIN CANCER	PRE-TEST CORRECT (%)	CORRECT (%) POST-TEST CORRECT (%)					
Basal cell carcinoma							
Picture A	49	97	<.001				
Picture B	43	100	<.001				
Squamous cell carcinoma							
Picture A	73	95	.01				
Picture B	67	100	.004				
Melanoma							
Picture A	0	97	<.001				
Picture B	59	97	.001				
N 27 fay and sammavisa							

Statistical significance (p) was determined using an exact version of the McNemar chi-square test

of skin cancer. Participants were then shown a series of 12 color images of noncancerous and cancerous cutaneous lesions on a tablet computer. For each image, subjects were asked to make a binary decision as to whether the lesion appeared to be malignant "cancer" or benign "not cancer." The 12 images consisted of two examples each of six common skin lesions (e.g., malignant melanoma, cherry angioma, seborrheic keratosis, benign melanocytic nevus, basal cell carcinoma, and squamous cell carcinoma). Afterwards, the participants were shown a two-minute educational video and surveyed again using the same 12 images. Preand postvideo scores were computed, with 12 points possible; one point was given for each correct answer. This Spanish video was scripted at the seventh-grade level, filmed with Hispanic healthcare providers speaking slowly, with accompanying captions to highlight key concepts. The video presented information about cancerous and noncancerous skin lesion detection, prevention, and screening.

RESULTS

A total of 37 subjects participated in the study (Table 1). Ninety-seven percent of participants reported having never seen a dermatologist. None had a personal history of skin cancer nor a family history of melanoma. Ninety-two percent admitted they never applied sunscreen during prolonged sun exposure. Of the eight percent who stated they used sunscreen, only five percent said they reapplied sunscreen when outdoors. Regarding educational format preference (video vs. written) 67.6 percent of the patients preferred video instruction. The video format preferences were not statistically significant. Pre-viewing baseline median score was six points (interquartile range [IQR]: 5–6 points). Postviewing median score improved to 11 points (IQR: 11–12 points), making the overall change from pre- to postintervention statistically significant (median=5 points, IQR: 4–6 points; p<.001). With regard to nonmelanoma skin cancers, squamous cell carcinoma identification improved from 74 percent to 98 percent (55/74 to 72/74), and basal cell identification rose from 35 percent to 99 percent (26/74 to 73/74). Initially, only 30 percent of subjects were able to identify melanoma prior to educational intervention. After watching the video, 97 percent could

BRIEF REPORT

identify this potentially deadly form of cancer (Table 2). A subject's education level and Fitzpatrick Skin Type did not correlate with score attained.

DISCUSSION

To date, little has been written about the impact or effectiveness of dermatology video education in the primary Spanish-speaking population of the US. The 2010 National Health Interview Survey revealed that only about seven percent of Hispanics reported ever having had a full-skin examination, compared to 25 percent of non-Hispanic Caucasian adults.8 This in part helps to explain why melanoma commonly presents in a more advanced stage with higher mortality among Hispanic individuals. Our educational video was effective in raising awareness about skin cancer and prevention methods while also providing participants with basic knowledge about visual features of common benign and malignant cutaneous lesions.

Limitations. Limitations of our study include the lack of a control group (i.e., a randomized design), not measuring the longterm retention of the information learned, and not characterizing the impact that viewing the video had on sun exposure and protection practices.

CONCLUSION

The use of video education in primary Spanish-speaking patients might be an

effective tool to enhance knowledge about skin health and improve identification of skin cancer. Additional research with larger patient numbers, control groups, and long-term outcome measures that include degree of information retention and the effects on skin protections practices are needed to confirm our findings.

SUPPLEMENTAL VIDEO

To view the educational video used in this study, please visit https://jcadonline.com/ supplemental-video-to-video-educationto-promote-skin-cancer-awareness-andidentification-in-spanish-speaking-patients/.

REFERENCES

- 1. Humes K, Jones N, Ramirez R. Overview of race and Hispanic origin: 2010. United States Census Bureau. March 2011. https://www.census.gov/ prod/cen2010/briefs/c2010br-02.pdf. Accessed 17 Jan 2020.
- Cress RD, Holly EA. Incidence of cutaneous 2. melanoma among non-Hispanic whites, Hispanics, Asians, and blacks: an analysis of california cancer registry data, 1988-93. Cancer Causes Control CCC. 1997;8(2):246-252.
- 3. Javed S, Javed SA, Mays RM, Tyring SK. Clinical characteristics and awareness of skin cancer in Hispanic patients. Dermatol Online J. 2013;19(9):19623.
- 4. Mahler HIM. Reasons for using and failing to use sunscreen: comparison among whites, Hispanics, and Asian/Pacific Islanders in Southern

- California. *JAMA Dermatol*. 2014;150(1):90-91. 5. Lenczowski E, Tung-Hahn E, Higareda J, et al. Video education to improve recognition of common benign and malignant cutaneous lesions and skin cancer prevention in the public. Int J Womens Dermatol. 2018;4(2):80-82.
- 6. Armstrong AW, Idriss NZ, Kim RH. Effects of video-based, online education on behavioral and knowledge outcomes in sunscreen use: a randomized controlled trial. Patient Educ Couns. 2011;83(2):273-277.
- 7. Trinh N, Novice K, Lekakh O, et al. Use of a brief educational video administered by a portable video device to improve skin cancer knowledge in the outpatient transplant population. Dermatol Surg Off Publ Am Soc Dermatol Surg Al. 2014;40(11):1233-1239.
- 8. Coups EJ, Stapleton JL, Hudson SV, et al. Skin cancer screening among Hispanic adults in the United States: results from the 2010 National Health Interview Survey. Arch Dermatol. 2012;148(7):861-863.
- 9. Harvey VM, Oldfield CW, Chen JT, Eschbach K. Melanoma disparities among US Hispanics: use of the Social Ecological Model to contextualize reasons for inequitable outcomes and frame a research agenda. J Skin Cancer. 2016;2016:4635740. JCAD